

BL Lac Objects and Relativistic Beaming Model

J.H. Fan

CRAL Observatoire de Lyon, 9 Avenue Charles Andr, 69 563 Saint-Genis-Laval Cedex,

France; and

Center for Astrophysics, Guangzhou Normal University, Guangzhou 510400, China

Received _____; accepted _____

ABSTRACT

In this paper: 1. The assumption of the dependence of Doppler factor on the emission frequency ($\delta_\nu \approx \delta_O^{1+\frac{1}{8}\log\frac{\nu}{\nu_o}}$, Fan et al. 1993) is used to explain the observational differences between the radio-selected BL Lac objects(RBLs) and the X-Ray-Selected BL Lac objects (XBLs): a) Hubble relation; b) different multiwavelength correlations; c) different regions in the effective spectral index ($\alpha_{RO} - \alpha_{OX}$) diagram; d) different polarization. The results suggest that RBLs and XBLs are the same. 2. From the analysis of the relation between infrared magnitude and redshift, it is proposed that the parent population of BL Lac objects should be FRI radio galaxies and FRII(G) radio galaxies showing the optical spectra of a galaxy. 3. From the superluminal motion, the assumption ($\delta_\nu \approx \delta_O^{1+\frac{1}{8}\log\frac{\nu}{\nu_o}}$) is confirmed. 4. Based on the relation between polarization and Doppler factor (Fan et al. 1997), it is proposed that the f , ratio of the beamed luminosity to the unbeamed luminosity in the source frame of OVV/HPQs is smaller than that of BL Lac objects: $f_{RBLs} \sim 6f_{FSRQs}$.

Subject headings: Beaming effect – BL Lacertae Objects (RBLs and XBLs) - OVV/HPQs – Superluminal Motion–Unified Scheme

1. Introduction

BL Lacertae objects are an extreme subclass of AGNs showing rapid and large amplitude variability, high and variable polarization, no or weak emission features ($EW < 5\text{\AA}$). From survey, BL Lacertae objects can be divided into radio selected BL Lac objects (RBLs) and X-ray selected BL Lac Objects (XBLs); or into quasar-like BL Lacertae objects (Q-BLs) and X-ray strong BL Lacertae objects (X-BLs) (Giommi et al. 1990) and into high frequency peaked BL Lac objects (HBLs) and low frequency peaked BL Lac objects (LBLs) (Padovani & Giommi 1995; Urry & Padovani 1995; Urry 1998).

There are some obvious differences between the two subclasses. Some authors claimed that they are the same class with XBLs having wider viewing angle than do RBLs (Ghisellini & Maraschi, 1989; Xie et al. 1993; Georganopoulos & Marsher 1996,1998; Fan et al. 1997a,b); Some authors claimed that there is a continuous spectral sequence from XBLs(HBLs) to RBLs(LBLs) (Sambruna et al. 1996).

2. Difference and Unity of BL Lac objects

2.1. The Observational Differences between RBLs and XBLs

- 1) RBLs do not fit the Hubble diagram so well as do XBLs;
- 2) RBLs do not show any multiwavelength correlation as do XBLs;
- 3) RBLs have higher radio and optical luminosities than do XBLs but they both have almost the same X-ray luminosity within the error range;
- 4) They occupy different regions in the effective spectral index diagram;
- 5) The averaged polarization of XBLs is lower than that of RBLs with $\overline{\Pi_{RBLs}} > 10\%$ and $\overline{\Pi_{XBLs}} < 5\%$.

2.2. Unity of RBLs and XBLs

The relativistic beaming model (Blandford & Rees 1978) suggests that the observed flux has been boosted. $S^{ob.} = \delta^p S^{in.}$ which gives

$$m_O^{in.} = m_O^{ob.} + 2.5 \times p \times \log \delta_O$$

, we think that should use the intrinsic data to compare the properties of RBLs and XBLs: For Hubble relation, We proposed a method to estimate δ_O (Xie et al. 1991). When we use the $m_V^{in.}$ to discuss the Hubble relation, much better results show up, and the corrected data of RBLs fit the same Hubble relation as XBLs (Xie et al. 1991; Fan et al. 1994) suggesting they are a single class with RBLs being more strongly beamed than are XBLs. From the microvariability amplitude, Miller & Noble (1996) also showed that RBLs have higher Lorentz factor than XBLs.

We assume that the Doppler factor satisfies (Fan et al. 1993):

$$\delta_\nu \approx \delta_O^{1 + \frac{1}{8} \log \frac{\nu}{\nu_O}}$$

which gives $\delta_R \sim \delta_O^{1.65}$. This is confirmed by the superluminal motion which gives $\delta_R \sim \delta_O^{1.93 \pm 0.22}$ (Fan et al. 1996a,b).

When the assumption is used on some RBLs with known optical Doppler factors, we found that the corrected luminosity of RBLs are equal to those of XBLs within the error range, the corrected RBLs data show good multiwavelength correlations, and the corrected spectral indices of RBLs shift to the XBLs region (see Fan et al. 1993, 1994; Fan & Xie 1996). The results suggest that RBLs are more strongly beamed than XBLs and that RBLs have smaller viewing angle than XBLs. So, we hope to use this assumption to explain the difference in their polarizations.

Following the work of Urry & Shafer (1984) the observed luminosity consists of two parts, the beamed and the unbeamed, $S_j^{in} = f S_{unb}$, $S^{ob} = (1 + f) S_{unb}$, and assuming

that the beamed flux consists of two proportional components (the polarized and the unpolarized ones),

$$S_{jp} = \eta S_{jup}, S_j^{in} = S_{jp} + S_{jup}$$

we obtained that the observed polarization can be written as

$$\Pi^{ob.} = \frac{(1+f)\delta^p}{1+f\delta^p}\Pi^{in.}$$

where, $\Pi^{in.} = \frac{f\eta}{(1+f)(1+\eta)}$. When we use the maximum polarization and the known Doppler factors (Fan et al. 1993; Ghisellini et al. 1993) to compare with the theoretical curve, The points fit the theoretical curves (see Fan et al. 1997a).

2.3. Parent Population of BL Lac Objects

Some authors think that the unified model includes BL Lac objects and FRI (Brown 1983; Urry et al. 1991; Bicknell 1994; Padovani 1998), others think that the unified model includes BL Lac objects, FRI and some FRII as well (Owen et al. 1996, see also Rector 1998). We think that the unified model should include BL Lac objects, FRI and FRII(G) radio galaxy with galaxy spectrum (Xie et al. 1993).

Recently, We used the faintest observed K magnitude to discuss the K-band Hubble relation and found that the corrected magnitude of RBLs, $K^{corr.}$, fit the same Hubble relation as the uncorrected X-ray selected BL Lac objects (XBLs), FRI, and FRII(G) confirming that BL lac objects, FRI and FRII(G) are unified (Fan et al. 1997b, see also Fig. 1).




Fig. 1.— The K-band Hubble diagram for BL Lac objects, FRI and FRII(G)

3. RBLs and Flat Spectral Radio Quasars (FSRQs)

The relation between RBLs and FSRQs (HPQs/OVV) is complex (Fan 1997a, Scarpa & Falomo 1997). We think FSRQs have smaller f than RBLs: From the core-dominance parameter (R) expression, $R = f\delta^p$, we found that FSRQs have smaller f 's than do RBLs (see Fan 1997b). From the polarization-Doppler factor relation we can see FSRQs lie in a region corresponding to smaller f 's as compared with RBLs (Fig. 2). So, we can get that FSRQs have smaller f than RBLs, and a value of $f_{RBLs} \sim 6f_{FSRQs}$ can be estimated from Fig. 2. It is clear that 3C279, 3C345 and 1156 are in the region of BL Lac objects. Does that mean that we should class them as RBLs? , or BL Lac objects and OVVs/HPQs are originally the same with BL Lac objects showing larger f .

I thank Prof. Roland Bacon, the director of l'Observatoire de Lyon for his supporting my attending the conference! This work is supported by the National Pandeng Project of China




Fig. 2.— Plot of Maximum polarization and Doppler factor for BL Lac objects and FSRQs

REFERENCES

- Bicknell G.V. 1994, ApJ, 422, 542
- Blandford R.D. & Rees M.J. 1978, in Pittsburgh Conf. on BL Lac, P328
- Brown, I.A.W. 1983, MNRAS, 204, 23p
- Fan J.H. et al. 1997a, A&A, 327, 947; 1997b, Ap&SS, 253, 275
- Fan J.H. 1997a, Ap.L. & Com. 35, 361; 1997b, in IAUS, 184, 242
- Fan J.H. et al. 1996a, A&AS, 116, 409; 1996b, IAUC. 159, p74
- Fan J.H. & Xie G.Z. 1996, A&A, 306, 55
- Fan J.H. et al. 1994a, A&AS, 105, 415; 1994b, Ap&SS, 213, 305
- Fan J.H. et al. 1993, ApJ, 415, 113
- Ghisellini, G. et al. 1993, ApJ, 407, 65
- Ghisellini, G. & Maraschi L. 1989, ApJ, 340, 181
- Giommi P. Barr, P. Pollock A.M.T. et al. 1990, ApJ, 356, 432
- Georganopoulos M & Marsher A.P. 1998, this conf.
- Georganopoulos M & Marsher A.P. 1996, ASP conf. Ser. Vol 110, P262
- Miller H.R. & Noble J.C. 1996, ASP Conf. Ser. Vol. 110, p17
- Owen F.N. et al. 1996, AJ, 111, 53
- Padovani P. & Giommi P. 1995, MNRAS 277 1477
- Rector, T.A. 1998, this conference

- Padovani P. 1998, this conference
- Sambruna R.M. Maraschi L. Urry C.M. 1996 ApJ, 463, 444
- Scarpa R. & Falomo R. 1997, A&A, 325, 109
- Urry C.M. 1998 this conference
- Urry C.M. & Padovani P. 1995, PASP, 107, 803
- Urry C.M. et al. 1991, ApJ, 382, 501
- Urry C.M. & Shafer R.A. 1984, ApJ, 280, 569
- Xie G.Z. et al. 1993, A&A, 278, 6
- Xie G.Z. et al. 1991, Ap&SS, 179, 321